

## Effectiveness of transcutaneous electrical nerve stimulation in management of neuropathic pain in patients with post traumatic incomplete spinal cord injuries

Amir Zeb<sup>1</sup>, Aatik Arsh<sup>2</sup>,  
Sher Bahadur<sup>3</sup>, Syed Muhammad Ilyas<sup>4</sup>

### ABSTRACT

**Objective:** To determine the effectiveness of Transcutaneous Electrical Nerve Stimulation (TENS) in management of neuropathic pain in post-traumatic incomplete spinal cord injury patients.

**Methods:** A quasi-experimental study was conducted from January 2017 to June 2017 at Paraplegic Center Hayatabad, Peshawar. Total 60 incomplete spinal cord injured patients with diagnosis of neuropathic pain were subjected to high frequency TENS of 80 HZ. One session was of 45 minutes while there were two sessions per day. TENS was applied for four days in a week and all patients were followed for eight week duration. Pain intensity was measured by using VAS (Visual analogue scale).

**Results:** Mean pain intensity on VAS at baseline was 6.45 which was decreased to 4.77 post intervention at day-1 while it was decreased to 3.48 at day-4 of week one. After application of TENS for 8 weeks, mean pain intensity was decreased to  $2.80 \pm 1.74$ . During the consecutive sessions of the TENS application, the pain intensity decreases in a linear fashion and there were significant difference ( $p < 0.05$ ) between pre and post treatment sessions.

**Conclusion:** TENS is useful and safe adjuvant in spinal cord injury patients for the management of neuropathic pain.

**KEYWORDS:** Neuropathic pain, Pain management, Physiotherapy, Spinal cord injury, TENS.

doi: <https://doi.org/10.12669/pjms.345.15659>

### How to cite this:

Zeb A, Arsh A, Bahadur S, Ilyas SM. Effectiveness of transcutaneous electrical nerve stimulation in management of neuropathic pain in patients with post traumatic incomplete spinal cord injuries. *Pak J Med Sci.* 2018;34(5):1177-1180.

doi: <https://doi.org/10.12669/pjms.345.15659>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Amir Zeb, PPDPT, MSPT.  
Senior Physical Therapist,
2. Aatik Arsh, DPT, MSPT.  
Physical Therapist,
3. Sher Bahadur, MPH.  
Senior Research Officer Rehman Medical College,
4. Syed Muhammad Ilyas, MSPT.  
Chief Executive Officer,

1,2,4: Paraplegic Center,  
Peshawar, Pakistan.

Correspondence

Dr. Aatik Arsh, DPT, MSPT.  
Physical Therapist,  
Paraplegic Center,  
Peshawar, Pakistan.  
E-mail: [aatikarshkmu@yahoo.com](mailto:aatikarshkmu@yahoo.com)

- \* Received for Publication: May 18, 2018
- \* 1st Revision Received: May 24, 2018
- \* 2nd Revision Received: September 5, 2018
- \* Final Revision Accepted: September 7, 2018

### INTRODUCTION

Spinal Cord Injury (SCI) is a devastating disability known to human kind.<sup>1</sup> Despite the fact that SCI is associated with incredible costs and human sufferings, yet exact statistics of SCI are not available in majority of developing countries including Pakistan.<sup>2</sup> The only available literature related to SCI in Pakistan is in the form of small single center based retrospective surveys.<sup>3-8</sup>

Pain is a common complication after SCI with prevalence of 18 to 96%, and almost 30% of these pain are diagnosed as neuropathic pain.<sup>9</sup> The intensity of the pain varies amongst these patients and it has been reported that 77.7% of patients with spinal cord injuries have moderate to severe pain.<sup>10</sup> Neuropathic pain mostly starts soon after injury

and can continue for the rest of the patient's life. The consequences of long term pain have often been associated negatively with the outcomes of rehabilitation.<sup>11</sup> Neuropathic pain is usually managed with a variety of pharmacological and non-pharmacological therapies<sup>12</sup> Among non-pharmacological treatments Transcranial Electrical Stimulation (TES), acupuncture, massage therapy and Transcutaneous Electrical Nerve Stimulation (TENS) are some of the frequently used modalities/ techniques for relieving neuropathic pain in SCI patients. TENS is one of the commonly used physical therapy modality for the management of pain in Pakistan.<sup>13-15</sup> In some clinical trial it was found that TENS had positive effects on management of pain in patients with SCI.<sup>10</sup> However, other studies on the efficacy of TENS in relieving neuropathic pain in SCI patients have not shown any effects and/or shows contradictory outcomes.<sup>10,16,17</sup> Due to controversy in literature, there was a dire need to conduct this study in order to determine the effectiveness of TENS in management of neuropathic pain in patients with post traumatic incomplete spinal cord injuries.

## METHODS

This quasi experimental study was conducted from January 2017 to June 2017 at Paraplegic Center Hayatabad, Peshawar. Paraplegic center Peshawar is second to none in the country providing comprehensive rehabilitative services to patients with spinal cord injuries.

A total of 60 patients with incomplete spinal cord injuries with diagnosis of neuropathic pain and age 20-60 years were recruited in the study using consecutive sampling technique. Complete spinal cord injury patients and/or those with complications e.g. pressure sores, fractures etc. were excluded.

TENS with high frequency of 80 HZ was applied to patients of SCI. Time for one session was 45 mints, while there were 2 sessions per day i.e. morning and evening session. TENS was applied for four consecutive days i.e. Monday, Tuesday, Wednesday, and Thursday. Each and every patient who entered into the study was followed for 8 weeks. Pain intensity was measured by using VAS before and after the application of TENS. Comparison between pre and post interventional scores was completed to see whether improvement occurred or not. Ethical approval was obtained from institutional ethical review committee of Paraplegic center, Peshawar.

## RESULTS

A total of 60 subjects with mean age  $52.64 \pm 0.48$  (ranged from 20-60 years) participated in the study, out of whom 75% (n=45) were male and 25% (n=15%) were female. Majority of the participants (71.7%, n=43) were married while the rest of participants (28.3%, n=17) were single. All the patients were from Khyber Pakhtunkhwa and were Pashto speaking. Most of the participants were laborer (55.0%, n=33), 13 (21.7%) participants were House wives while rest of the participants (23.3%, n=14) were having other professions. Forty six (76.7%) participants had thoracic paraplegia, 11 (18.3%) patients had lumbar paraplegia while 3 (5.0%) patients had cervical tetraplegia.

Mean pain intensity on VAS at baseline was  $6.45 \pm 1.09$  which were decreased to  $4.77 \pm 1.52$  post intervention at day-1 of week 1. The mean decreased in the pain intensity was seen through the array of intervention. The mean pain intensity was decreased to  $3.48 \pm 1.91$  at day-4 of week-1. During the consecutive sessions of TENS application, the pain intensity decreases in a linear fashion and there were significant difference ( $p < 0.05$ ) between pre and post treatment sessions. (Table-I)

After application of TENS for 8 weeks, mean pain intensity at day 1 of week 8 was  $2.99 \pm 1.33$  which was decreased to  $2.92 \pm 1.2$  on day-two and  $2.91 \pm 1.5$  on day three while  $2.80 \pm 1.7$  on day-4. There was significance difference ( $p < 0.05$ ) in pain intensity at pre and post intervention of each session. (Table-I)

Table-I: Comparison of Pain at pre and post treatment at different days of Week 1 and Week 8.

Pain at pre & post treatment		Mean $\pm$ SD	P-Value
Week 1, Day-1,	D1Pre	$6.45 \pm 1.09$	<0.001
	D1Post	$4.77 \pm 1.52$	
Week 1, Day-2	D2Pre	$5.96 \pm 1.20$	< 0.001
	D2Post	$4.06 \pm 1.57$	
Week 1, Day-3	D3Pre	$5.63 \pm 1.39$	< 0.001
	D3post	$3.88 \pm 1.99$	
Week 1, Day-4	D4pre	$5.37 \pm 2.00$	< 0.001
	D4post	$3.48 \pm 1.91$	
Week 8, Day-1,	D1Pre	$4.21 \pm 1.78$	<0.001
	D1Post	$2.99 \pm 1.33$	
Week 8, Day-2	D2Pre	$3.92 \pm 1.69$	< 0.001
	D2Post	$2.92 \pm 1.27$	
Week 8, Day-3	D3Pre	$3.41 \pm 1.52$	< 0.001
	D3post	$2.91 \pm 1.52$	
Week 8, Day-4	D4pre	$3.10 \pm 1.42$	< 0.001
	D4post	$2.80 \pm 1.74$	

## DISCUSSION

SCI is associated with various degree of sensory and motor dysfunctions.<sup>18</sup> Neuropathic pain is common among spinal cord injury patients which can be managed both through pharmacological as well as Non-Pharmacological interventions.

Using the Quasi experimental design this study was aimed to determine the effectiveness of TENS in management of neuropathic pain in patients with post traumatic incomplete SCI patients. Result indicated that mean decreased in the pain intensity was seen through the array of intervention. During the consecutive sessions of the physical therapy the pain intensity decreased in a linear fashion and there are significant difference ( $p < 0.05$ ) between pre and post treatment session. Following TENS in patients with post Traumatic incomplete spinal cord injuries, mean pain score at week week-8 and day-4 was decreased to 2.80 which reveals a marked decreased in pain intensity. These results are consistent with a randomized control study which reveals that there was a marked decreased in the pain intensity after each consecutive intervention (application of TENS).<sup>17</sup> Results indicates that after each session the pain decreases to a significant level ( $p < 0.05$ ). The results of the present study are in consistence with studies conducted by Norrbrink C et al. and Celik EC et al. who found that TENS complement pharmacological treatment in patients with SCI and neuropathic pain.<sup>9,17</sup> From present study it was meticulous to see the low pain score (3.78) at base line of week-8, as it was 6.45 at week-1 and day-1. Similar finding was also reported by Kilinc M et al. who compared pre- and post TENS treatment for patients with peripheral (PNP) or central neuropathic pain (CNP). At the beginning of the trial, the minimal, maximal, mean, and current pain intensities were similar between the CNP and PNP groups. Post-treatment pain intensity values were significantly lower than pre-treatment values in both groups ( $p < 0.05$ ).<sup>19-20</sup>

Certain adverse affects of TENS has been reported in literature.<sup>17</sup> For example, high intensity TENS can cause rashes and local tingling sensations.<sup>20</sup> However these adverse reactions are minimal as compared to its beneficial effects. Though in current study patients were followed for 8 weeks, however there some studies reported relapse of neuropathic pain in spinal cord injury patients.<sup>21-23</sup> There is possibility of relapse of neuropathic pain in participants of current study, therefore proper long term follow up is advised to all participants.

TENS is primarily aimed to provide a degree of symptomatic pain relief by exciting sensory nerves.<sup>21-22</sup> The effectiveness of TENS varies with the clinical pain being treated, but research suggest that when used 'well' it provides significantly greater pain relief than a placebo intervention.<sup>23</sup>

Despite the fact that current study is first of its kind conducted in Pakistan which reported effectiveness of TENS in managing neuropathic pain in spinal cord injury patients, however it has some limitations. First of all, its design was quasi-experimental so the results of current study can be affected by confounding variable. Secondly sample size of current study was small due to which generalizability of the results of current study is questionable.

## CONCLUSION

TENS is useful and safe adjuvant in spinal cord injury patients for the management of neuropathic pain. A consistence and long term rehabilitation with TENS in patients with post Traumatic incomplete SCI is useful.

*Grant Support & Financial Disclosures:* None.

*Declaration:* The results of current study was presented at 5th International Conference and Expo on Novel Physiotherapies J Nov Physiother 2018, Volume 8. March 19-20, 2018 | Berlin, Germany. "Effectiveness of transcutaneous electrical nerve stimulation in management of neuropathic pain in patients with post traumatic incomplete spinal cord injuries, at paraplegic center, Peshawar, Pakistan" Amir Zeb Parpalegic Center, Pakistan.

## REFERENCES

1. Lee B, Cripps R, Fitzharris M, Wing P. The global map for traumatic spinal cord injury epidemiology: update 2011, global incidence rate. *Spinal Cord*. 2014;52(2):110-106. doi:10.1038/sc.2012.158
2. Rathore FA, Farooq F, Muzammil S, New PW, Ahmad N, Haig AJ. Spinal cord injury management and rehabilitation: highlights and shortcomings from the 2005 earthquake in Pakistan. *Arch Phys Med Rehabil*. 2008;89(3):579-585. doi: 10.1016/j.apmr.2007.09.027.
3. Arsh A, Darain H, Haq ZU, Zeb A, Ali I, Ilyas SM. Epidemiology of spinal cord injuries due to bomb blast attacks, managed at paraplegic centre peshawar, pakistan: a nine years retrospective study. *Khyber Med Uni J*. 2017;9(2):55-59.
4. Arsh A, Darain H, Ilyas SM. Epidemiology and prevalence of deep venous thrombosis in patients with spinal cord injury. *Pak J Neurolog Sci*. 2016;11(4):22-25.
5. Arsh A, Darain H, Ilyas SM, Zeb A. Prevelacne of pressure ulcers in patients with spinal cord injury; a retrospective study. *Pak J Neurolog Sci*. 2012;11(4):16-21.

6. Arsh A, Darain H, Ilyas SM, Zeb A. Consequences of traumatic spinal cord injury during pregnancy in Pakistan. *Spinal Cord Series Cases*. 2017;3:17041. doi: 10.1038/scsanc.2017.41.
7. Arsh A, Darain H, Muhammad D, Khan MN, Ilyas SM, Zeb A, et al. FACTORS responsible for readmissions to rehabilitation center following spinal cord injury. *J Khyber Coll Dent*. 2017;7(2):60-65.
8. Darain H, Arsh A, Zeb A, Ilyas SM, Muhammad D, Khan MN. Epidemiology, clinical features and consequences of spinal cord injury in children. *J Coll Physicians Surg Pak*. 2018;28(7):532-535. doi: 10.29271/jcsp.2018.07.532
9. Celik E, Erhan B, Gunduz B, Lakse E. The effect of low-frequency TENS in the treatment of neuropathic pain in patients with spinal cord injury. *Spinal Cord*. 2013;51(4):334-337. doi: 10.1038/sc.2012.159. Repeated on Ref. 20
10. Sadeghi H, Banitalebi E, Raeisi Dehkordi M. The effect of body-weight-supported training exercises on functional ambulation profile in patients with paraplegic spinal cord injury. *Physical Treatments-Specific Phys Ther J*. 2015;4(4):205-212.
11. Werhagen L, Budh C, Hultling C, Molander C. Neuropathic pain after traumatic spinal cord injury—relations to gender, spinal level, completeness, and age at the time of injury. *Spinal Cord*. 2004;42(12):665-673. doi:10.1038/sj.sc.3101641
12. Rathore MFA, Hanif S, Farooq F, Ahmad N, Mansoor SN. Traumatic spinal cord injuries at a tertiary care rehabilitation institute in Pakistan. *J Pak Med Assoc*. 2008;58(2):53-57.
13. Arsh A, Darain H. Physical therapy modalities used by physical therapists in the management of chronic low back pain in khyber pakhtunkhwa. *South Asian J Med*. 2016;1(2):35-40.
14. Osama M. Use of Trans-cutaneous Electric Nerve Stimulation (TENS) can alleviate labour pain and delay the use of neuraxial analgesia. *J Pak Med Assoc*. 2018;68(4):681.
15. Yameen F, Shahbaz NN, Hasan Y, Fauz R, Abdullah M. Efficacy of transcutaneous electrical nerve stimulation and its different modes in patients with trigeminal neuralgia. *J Pak Med Assoc*. 2011;61(5):437-439.
16. Picavet H, Schouten J. Musculoskeletal pain in the Netherlands: prevalences, consequences and risk groups, the DMC 3-study. *Pain*. 2003;102(1):167-178. doi: 10.1016/s0304-3959(02)00372-x
17. Norrbrink C. Transcutaneous electrical nerve stimulation for treatment of spinal cord injury neuropathic pain. *J Rehabil Res Dev*. 2009;46(1):85-93. doi: 10.1682/JRRD.2008.04.0051. Repeated on Ref. 19
18. Hagen EM, Rekind T. Management of Neuropathic Pain Associated with Spinal Cord Injury. *Pain Ther*. 2015;4(1):51-65. doi: 10.1007/s40122-015-0033-y.
19. Arsh A, Jan A. Prevalence of low back pain among dpt students in Peshawar. *South Asian J Med*. 2015;1(2):29-34.
20. Kilinc M, Livanelioglu A, Yildirim SA, Tan E. Effects of transcutaneous electrical nerve stimulation in patients with peripheral and central neuropathic pain. *J Rehabil Med*. 2014;46(5):454-460. doi: 10.2340/16501977-1271.
21. Ozkul C, Kilinc M, Yildirim SA, Topcuoglu EY, Akyuz M. Effects of visual illusion and transcutaneous electrical nerve stimulation on neuropathic pain in patients with spinal cord injury: A randomised controlled cross-over trial. *J Back Musculoskelet Rehabil*. 2014;28(4):709-719. doi: 10.3233/BMR-140573.
22. Vugt MAP, Joosen MCW, Mert A, Zedlitz A, Vrijhoef HJM. Serious gaming during multidisciplinary rehabilitation for patients with complex chronic pain or fatigue complaints: study protocol for a controlled trial and process evaluation. *BMJ Open*. 2017;7(6):e016394. doi: 10.1136/bmjopen-2017-016394.
23. Sivaramakrishnan A, Solomon JM, Manikandan N. Comparison of transcutaneous electrical nerve stimulation (TENS) and functional electrical stimulation (FES) for spasticity in spinal cord injury - A pilot randomized cross-over trial. *J Spinal Cord Med*. 2017;1-15. doi: 10.1080/10790268.2017.1390930.

#### ***Authors' Contribution:***

**AZ:** Concept and study design, literature search and literature review, Acquisition of data.

**AA:** Acquisition of data, drafting the manuscript.

**SB:** Analysis & interpretation of data.

**SM:** Critical revision, final approval of the version to be published.